

## 2110

**Breeding diet for rabbits**

The 2110 formula is a cereal- (soy, wheat, corn) and alfalfa-based fixed formula which is free of fish/animal meal and deficient in nitrosamines. This long-term breeding diet is rich in crude fibre and was designed as complete feedingstuff for breeding adult rabbits and young rabbits up to an age of 10 weeks. Addition of hay is recommended.

**Available as**

2113 - 4mm pellets

**Product variations**

The 2110 formula is optionally also available in one or a combination of the following variants.

**Variant Fortified**

autoclavable, increased vitamin content, coated on request

**Variant Irradiated**

irradiated with 5 to 50kGy

*The contents may vary for these variants.*

**Metabolized energy**

| <b>Content</b> | <b>Value</b> | <b>unit</b> |
|----------------|--------------|-------------|
| Fat            | 319 (12%)    | kcal/kg     |
| Protein        | 611 (24%)    | kcal/kg     |
| Carbonhydrates | 1,663 (64%)  | kcal/kg     |

**crude nutrients and moisture**

| <b>Content</b>           | <b>Value</b>    | <b>unit</b> |
|--------------------------|-----------------|-------------|
| Moisture                 | 113,622 (11.4%) | mg/kg       |
| Crude Ash                | 75,567 (7.6%)   | mg/kg       |
| Crude Fibre              | 195,168 (19.5%) | mg/kg       |
| Crude Fat                | 35,417 (3.5%)   | mg/kg       |
| Crude Protein            | 152,842 (15.3%) | mg/kg       |
| Nitrogenfree extractives | 427,384 (42.7%) | mg/kg       |

## Carbonhydrates

| Content         | Value   | unit  |
|-----------------|---------|-------|
| Monosaccharides | 0       | mg/kg |
| Disaccharides   | 33,878  | mg/kg |
| Polysaccharides | 298,293 | mg/kg |

## Minerals

| Content    | Value  | unit  |
|------------|--------|-------|
| Calcium    | 10,642 | mg/kg |
| Potassium  | 8,940  | mg/kg |
| Magnesium  | 1,513  | mg/kg |
| Sodium     | 2,481  | mg/kg |
| Phosphorus | 5,147  | mg/kg |

## Trace elements

| Content    | Value    | unit  |
|------------|----------|-------|
| Aluminium  | 326.24   | mg/kg |
| Chlorine   | 4,455.60 | mg/kg |
| Iron       | 181.23   | mg/kg |
| Flourine   | 4.83     | mg/kg |
| Iodine     | 1.38     | mg/kg |
| Cobalt     | 0.33     | mg/kg |
| Copper     | 11.16    | mg/kg |
| Manganese  | 62.90    | mg/kg |
| Molybdenum | 0.48     | mg/kg |
| Sulfur     | 1,994.70 | mg/kg |
| Selenium   | 0.23     | mg/kg |
| Zinc       | 56.29    | mg/kg |

## Added vitamins

| Content          | Value  | unit  |
|------------------|--------|-------|
| Vitamin A        | 15,000 | IU/kg |
| Vitamin D3       | 600    | IU/kg |
| Vitamin E        | 76     | mg/kg |
| Vitamin K3       | 3      | mg/kg |
| Vitamin B1       | 18     | mg/kg |
| Vitamin B2       | 12     | mg/kg |
| Vitamin B6       | 9      | mg/kg |
| Vitamin B12      | 24     | µg/kg |
| Nicotinic acid   | 36     | mg/kg |
| Pantothenic acid | 21     | mg/kg |
| Folic acid       | 4      | mg/kg |
| Biotin           | 375    | µg/kg |
| Choline chloride | 984    | mg/kg |
| Vitamin C        | 36     | mg/kg |

## Amino acids

| Content       | Value  | unit  |
|---------------|--------|-------|
| Alanine       | 5,539  | mg/kg |
| Arginine      | 6,778  | mg/kg |
| Aspartic acid | 9,962  | mg/kg |
| Cystine       | 1,713  | mg/kg |
| Glutamic acid | 20,234 | mg/kg |
| Glycine       | 5,120  | mg/kg |
| Histidine     | 2,524  | mg/kg |
| Isoleucine    | 4,551  | mg/kg |
| Leucine       | 8,054  | mg/kg |
| Lysine        | 5,033  | mg/kg |
| Methionine    | 1,623  | mg/kg |
| Phenylalanine | 5,022  | mg/kg |
| Proline       | 6,760  | mg/kg |
| Serine        | 5,301  | mg/kg |
| Threonine     | 4,198  | mg/kg |
| Tryptophan    | 1,533  | mg/kg |
| Tyrosine      | 3,611  | mg/kg |
| Valine        | 5,444  | mg/kg |

## Fatty acid

| Content                     | Value | unit  |
|-----------------------------|-------|-------|
| Arachidic acid C-20:0       | 44    | mg/kg |
| Eicosanoic acid C-20:1      | 55    | mg/kg |
| Alpha-Linolenic acid C-18:3 | 1,233 | mg/kg |
| Linolenic acid C-18:2       | 8,998 | mg/kg |
| Palmitic acid C-16:0        | 2,175 | mg/kg |
| Stearic acid C-18:0         | 661   | mg/kg |
| Oleic acid C-18:1           | 3,800 | mg/kg |